

On magnetic field generation in Kolmogorov turbulence

Stanislav Boldyrev¹ and Fausto Cattaneo²

¹*Department of Astronomy and Astrophysics, University of Chicago, 5640 S. Ellis Ave, Chicago, IL 60637*

²*Department of Mathematics, University of Chicago, 5734 S. University, Chicago, IL 60637*

(February 2, 2008)

We analyze the initial, kinematic stage of magnetic field evolution in an isotropic and homogeneous turbulent conducting fluid with a rough velocity field, $v(l) \sim l^\alpha$, $\alpha < 1$. We propose that in the limit of small magnetic Prandtl number, i.e. when ohmic resistivity is much larger than viscosity, the smaller the roughness exponent α , the larger the magnetic Reynolds number that is needed to excite magnetic fluctuations. This implies that numerical or experimental investigations of magnetohydrodynamic turbulence with small Prandtl numbers need to achieve extremely high resolution in order to describe magnetic phenomena adequately.

1. Introduction. In a turbulent highly conducting fluid, magnetic fields may be amplified since the field lines are generally stretched by randomly moving fluid elements in which these lines are frozen [1]. Such a mechanism of turbulent dynamo is expected to work in a variety of astrophysical systems (galaxy clusters, interstellar medium, stars, planets), is confirmed numerically, and is consistent with simple analytical models.

Valuable insight can be gained from considering the so-called kinematic stage of the dynamo, when magnetic field is amplified from an initially weak ‘seed’ field. As the simplest example, assume that the resistive scale is smaller than or of the same order as the viscous scale of the fluid. In the turbulent Kolmogorov velocity field, the smallest eddies have the highest shearing rate given by $v(l)/l \sim l^{-2/3}$, where l is the size of the eddy, and $v(l) \sim l^{1/3}$ is its turn-over velocity. Therefore, at this stage the magnetic field grows predominantly on small scales. Fig. 1 shows schematically the initial stage of the evolution of the magnetic spectrum, see, e.g., [2].

The magnetic energy collapses toward small, resistive scale during this initial evolution, until the field is strong enough to affect the dynamics of fluid through the Lorentz force. Such behavior is the evidence of small-scale turbulent dynamo; it is firmly established in numerical experiments, and can be derived analytically [2,3,5–9]. Since in this example, the resistive scale is smaller than the viscous one, the dynamo is essentially governed by a smooth, viscous-scale velocity field.

To study the growth of magnetic energy on larger scales we have to understand how the magnetic field is generated in the inertial interval of the turbulence (the interval of scales much smaller than the external scale where the turbulence is excited, and much larger than the viscous scale where the turbulent energy is dissipated). This question is non-trivial since in this interval the velocity field is not smooth, i.e., $v(l) \sim l^\alpha$, with $\alpha < 1$. This situation is especially relevant for the case of small magnetic Prandtl number ($\text{Pm} = \nu/\eta$, where ν is fluid viscosity and η is resistivity), where the magnetic energy is concentrated mostly in the inertial interval of the velocity field.

The question was first addressed by Batchelor [3] who

used the analogy between magnetic field lines and fluid vorticity lines to conclude that when $\nu \leq \eta$, magnetic energy is not amplified. This analogy was criticized in [4] because the magnetic field can have arbitrary initial conditions, while the vorticity field is related to the velocity field. It was further argued in [10] that magnetic field line stretching generally dominates resistive reconnection, thus making dynamo possible for $\nu \leq \eta$. Direct numerical simulations of MHD turbulence with $\text{Pm} = 1$ confirm that weak magnetic fluctuations are generally amplified by the Navier-Stokes velocity field except for a special case when the initial magnetic field is close to the vorticity field [11,12].




FIG. 1. Sketch of the initial stage of magnetic field amplification by turbulent velocity field. E_k is the spectrum of the velocity field in k space, H_k is the spectrum of the magnetic field. The viscous cut-off k_ν is larger than the resistive cut-off k_η . Magnetic field energy is shown for three consecutive times, it grows and gets accumulated at the resistive scale.

Recently, there appeared the number of numerical simulations of MHD turbulence with small magnetic Prandtl numbers, where magnetic fluctuations were not amplified [13–16], which revived the claims that dynamo does not exist in the Kolmogorov turbulence with $\text{Pm} \ll 1$, see e.g., [15,16]. However, astrophysical observations

show that magnetic fields are generated by turbulent motion rather effectively in planets and stars where magnetic Prandtl numbers are small (e.g., in the geodynamo, $\text{Pm} \sim 10^{-5} - 10^{-6}$, in the solar photosphere, $\text{Pm} \sim 10^{-7}$). This apparent contradiction motivated our interest to the problem.

In this paper we argue that dynamo always exists in a rough velocity field, such as the Kolmogorov field. We however find that the magnetic Reynolds number, $\text{Rm} = v(L_0)L_0/\eta$, and the numerical resolution required to generate magnetic field strongly depend on the roughness exponent of the velocity field, α ; the rougher the velocity field, the larger the required resolution. This result explains why dynamo action is hard to achieve in experiments with small magnetic Prandtl numbers, while it is easily achieved when the magnetic Prandtl number is large, $\text{Pm} \geq 1$. In the latter case the velocity field is effectively smooth at the resistive scales, where the magnetic energy is concentrated.

Our analysis is based on the so-called Kazantsev model of kinematic dynamo, which is exactly solvable and which allows us to change the velocity roughness exponent, α .

2. The Kazantsev model for a rough velocity field. Consider the induction equation for the magnetic field:

$$\partial B^i / \partial t + v^j \partial B^i / \partial x^j - B^j \partial v^i / \partial x^j = \eta \Delta B^i, \quad (1)$$

where $B^i(\mathbf{x}, t)$ is the magnetic field, $v^i(\mathbf{x}, t)$ is the fluid velocity, η is small ohmic resistivity, and we sum over the repeated vector indices. At the initial stage of evolution, the weak magnetic field is passively advected by the fluid. Therefore, one can use simplifying assumptions about the fluid velocity to make the problem manageable. Kazantsev [5] and Kraichnan [17] introduced the model based on the Gaussian, short-time correlated velocity field, with zero mean and the covariance

$$\langle v^i(\mathbf{x}, t) v^j(\mathbf{x}', t') \rangle = \kappa^{ij}(\mathbf{x} - \mathbf{x}') \delta(t - t'). \quad (2)$$

This model is a valuable tool for the analytical investigation of kinematic dynamo; direct numerical simulations reveal that a purely Gaussian velocity field amplifies small magnetic fluctuations in a similar manner as the true Navier-Stokes field [11].

Assuming isotropy and homogeneity, the velocity correlation function has the form

$$\kappa^{ij}(\mathbf{r}) = \kappa_N(r) \left(\delta^{ij} - \frac{r^i r^j}{r^2} \right) + \kappa_L(r) \frac{r^i r^j}{r^2}, \quad (3)$$

where $\mathbf{r} = \mathbf{x} - \mathbf{x}'$. If we further assume that the velocity field is incompressible, we have $\kappa_N = \kappa_L + (r \kappa'_L)/2$, and velocity statistics can be characterized by only one scalar function, $\kappa_L(r)$.

The model defined by (1), (2), and (3) allows one to write a closed equation for the correlation function of the magnetic field,

$$\langle B^i(\mathbf{x}, t) B^j(\mathbf{x}', t) \rangle = H^{ij}(\mathbf{x} - \mathbf{x}', t), \quad (4)$$

where, analogously to (3), the H^{ij} function can be represented as

$$H^{ij} = H_N(r, t) \left(\delta^{ij} - \frac{r^i r^j}{r^2} \right) + H_L(r, t) \frac{r^i r^j}{r^2}, \quad (5)$$

furthermore, the condition $\nabla \cdot \mathbf{B} = 0$ gives $H_N = H_L + (r H'_L)/2$. We will characterize the magnetic field correlator by the function $H_L(r, t)$. The equation for this function can be found by differentiating (4) with respect to time, and by using Eqs. (1)-(3). A rather tedious but essentially straightforward calculation gives:

$$\partial_t H_L = \kappa H''_L + \left(\frac{4}{r} \kappa + \kappa' \right) H'_L + \left(\kappa'' + \frac{4}{r} \kappa' \right) H_L, \quad (6)$$

where primes denote the derivatives with respect to r and we have introduced the ‘renormalized’ velocity correlation function $\kappa(r) = 2\eta + \kappa_L(0) - \kappa_L(r)$. Equation (6) was originally derived by Kazantsev [5], and can be rewritten in a different form that formally coincides with the Shrödinger equation in the imaginary time. Effecting the change of variable, $H_L = \psi(r, t) r^{-2} \kappa(r)^{-1/2}$, one obtains:

$$\partial_t \psi = \kappa(r) \psi'' - V(r) \psi, \quad (7)$$

which describes the wave function of a quantum particle with the variable mass, $m(r) = 1/[2\kappa(r)]$, in a one-dimensional potential ($r > 0$):

$$V(r) = \frac{2}{r^2} \kappa(r) - \frac{1}{2} \kappa''(r) - \frac{2}{r} \kappa'(r) - \frac{(\kappa'(r))^2}{4\kappa(r)}. \quad (8)$$

This equation can be investigated for different choices of $\kappa(r)$, however, we restrict ourselves to the inertial interval of the turbulence, where the velocity correlator has power-law asymptotics, $\kappa(r) \propto r^\beta$. The exponent β can be found from the scaling of turbulent diffusivity, $D \sim v(r)r \sim r^{1+\alpha}$. Indeed, in the derivation of Eq. (6) we used the integral $D = \int_0^\infty \langle [v(x, t) - v(x', t)][v(x, t') - v(x', t')] \rangle d(t - t') = \kappa(r)$, which is the turbulent diffusivity [10]. Comparing the two expressions we obtain $\beta = 1 + \alpha$.

The Shrödinger equation (7) has the effective potential $U_{eff}(r) = V(r)/\kappa(r) = A(\beta)/r^2$, where $A(\beta) = 2 - 3\beta/2 - 3\beta^2/4$. When $A(\beta) < -1/4$, the quantum particle falls toward the origin [18] and its wave function is therefore concentrated at the smallest, resistive scale. This behavior is the manifestation of the dynamo mechanism that we discussed in the introduction. We obtain that $A(1 + \alpha) < -1/4$ for any roughness exponent of the velocity field, $0 < \alpha < 1$, therefore, dynamo is always possible; the same conclusion was made in [10].

3. Magnetic field correlator and dynamo growth rates. To find the wave function (the magnetic field correlator) we need to specify the boundary conditions. The small-scale regularization is naturally given by ohmic resistivity. For scales much smaller than the correlation scale of the velocity field, the κ function can be expanded as

$\kappa(r) \simeq \kappa_L(0)(2\eta + r^{1+\alpha})$, which corresponds to the limit of infinitely small Prandtl number. In this formula and below we use the dimensionless variables: η is measured in the units of the large-scale turbulent diffusivity, $\kappa_L(0)$, while r is in the units of the integral scale, L_0 .

The boundary condition at the origin follows from finiteness of magnetic energy, $\lim_{r \rightarrow 0} H_L(r, t) = H_0(t) < \infty$. The boundary condition at large scales follows from the absence of the mean magnetic field, $\lim_{r \rightarrow \infty} H_L(r, t) = 0$. We will see in a moment that at the kinematic stage magnetic energy is concentrated at the resistive scale, and the corresponding wave function decays exponentially fast in the inertial interval, $r \gg r_\eta$, so it is indeed independent of the large-scale properties of the velocity field, as we expected.

Problem (6) can be cast into the Sturm–Liouville form by changing $H_L(r, t) = h(r, t)/r^2$, which teaches us that its solution can be expanded in the eigenfunctions of the corresponding Sturm–Liouville operator. The maximum growth rate is therefore given by the largest eigenvalue of the operator on the right-hand side of Eqs. (6) or (7). The corresponding wave function is the ground state of the potential $U_{eff}(r)$. Thus, following Kazantsev [5], we look for the solution of (7) in the form $\psi(r, t) = \varphi(r) \exp(\lambda t)$.

In the inertial interval, $2\eta \ll r^\beta$, the equation for the φ function reads

$$-\varphi'' + \left[\frac{-3\beta^2 - 6\beta + 8}{4r^2} + \frac{\lambda}{r^\beta} \right] \varphi = 0, \quad (9)$$

where λ is dimensionless and is measured in the units of $\kappa_L(0)/L_0^2$. For small λ , the potential is dominated by the first term in the square brackets, and one can easily check that the corresponding wave function oscillates in the inertial interval. However, the ground-state wave function cannot oscillate, therefore the ground-state growth rate, λ , will be such that the second term in the square brackets of (9) dominate the first one in the inertial interval. At the resistive scale, $r_\eta = (2\eta)^{1/\beta}$, both terms should be of the same order, therefore, $\lambda \sim \eta^{(\beta-2)/\beta}$.

The wave function corresponding to the growing solution, $\lambda > 0$, decays exponentially fast for $r \gg \eta^{1/\beta}$, $\varphi \propto \exp\left(-\frac{2\sqrt{\lambda}r^{(2-\beta)/2}}{2-\beta}\right)$. This function is concentrated at the resistive scale and its growth rate is of the order of the eddy turn-over time at this scale; in agreement with our qualitative discussion in the introduction. When one changes the magnetic Prandtl number (by changing viscosity, for instance) the effective roughness of the velocity field at the resistive scale changes. We therefore suggest that the effect of different Prandtl numbers can be studied in terms of Eq. (9) with different roughness exponents. By this analogy, the limit of the smooth velocity field, $\beta = 2$, corresponds to large Pm, while the other limit of the Kolmogorov-scaled velocity field, $\beta = 4/3$, corresponds to small Pm.

We observe that the rougher the velocity field, the broader the wave function compared to the resistive scale,

and, therefore, the larger the magnetic Reynolds number necessary to generate magnetic fluctuations. To estimate the critical resolution for the dynamo onset, we solved Eq. (6) numerically with the large-scale boundary condition, $H_L(\ell) = 0$. For given β and η , we increased the ‘system size’, ℓ , until H_L started to grow and we thus found the critical value $\ell = \ell_c$. An analogous calculation could be done by fixing ℓ and changing η .

To characterize the inertial range, we introduce the resolution parameter, $R = \ell_c/r_\eta = \ell_c/(2\eta)^{1/\beta}$; this parameter is universal, i.e., it is independent of large and small-scale properties of the velocity field. We obtain that when the velocity roughness increases from $\beta = 2.0$ to $\beta = 1.3$, the corresponding resolution parameter increases from $R \approx 3.8$ to $R \approx 29$, see Fig. (2). This means that if we simulated the random equation (1) directly, the required numerical resolution should increase by about an order of magnitude in each spatial direction as we go from the smooth to the Kolmogorov-scaled velocity field.




FIG. 2. The numerically found resolution parameter, $R = \ell_c/r_\eta$, as a function of the velocity roughness exponent, β .

The subresistive scales, $r \ll \eta^{1/\beta}$, can be easily described using Eq. (6). By direct substitution, one can check that the magnetic field correlator can be expanded as $H_L(r, t) = H_0[1 - r^\beta/(2\eta) + \dots] \exp(\lambda t)$, which implies that the spectrum of the magnetic field decays as $H_k \sim k^2 |B_k|^2 \propto k^{-1-\beta}$ in the subresistive region, $\eta^{-1/\beta} \ll k \ll \nu^{-1/\beta}$. With such a spectrum, the rate of magnetic energy dissipation at the viscous scale is bigger than the rate at the resistive scale by the factor $k_\eta^5 |B_{k_\eta}|^2 / k_\nu^5 |B_{k_\nu}|^2 \sim \text{Pm}^{-(2-\beta)/\beta}$. Thus, the effective resistivity may be ‘anomalously large’ in turbulent flows with small Prandtl numbers.

Strictly speaking, the last result requires short-time correlated small-scale eddies and is not applicable in the Kolmogorov turbulence where velocity is correlated at the eddy turn-over time, $\tau_{corr}(l) \sim l/v(l)$. At the subresistive scales, $l < l_\eta$, this correlation time is larger

than the resistive relaxation time, $\tau_\eta \sim l^2/\eta$, while in the Kazantsev model the velocity correlation time is smaller than the resistive relaxation time. However, model (2) is physically self-consistent. The large magnetic energy dissipation is balanced by the large energy transfer from the small-scale velocity eddies to the small scale magnetic field; this is possible since the fluid energy is formally infinite in (2). In a practical situation where the velocity correlation time is not infinitely small, the asymptotic tail $H_k \sim k^{-1-\beta}$ will hold up to the scales where this correlation time is comparable to the resistive time, $k \sim 1/\sqrt{\eta\tau_{corr}}$. One can expect that for smaller scales, the magnetic energy spectrum will have a steeper decay, see, e.g., [4,19,20].

4. Conclusions. We proposed that magnetic fluctuations are always generated in isotropic and homogeneous three-dimensional turbulence if the magnetic Reynolds number is large enough. The required critical magnetic Reynolds number sharply increases with velocity roughness, which explains the ‘no-dynamo’ outcomes reported in numerical simulations with small Prandtl numbers, see e.g., [13–16]. We do not see any physical reason for the absence of dynamo in these numerical simulations rather than lack of resolution. Our results also suggest that obtaining small-scale dynamo will be a serious challenge for laboratory experiments, where the magnetic Prandtl number is small, $Pm \sim 10^{-5}$, while the magnetic Reynolds number is rather moderate, $Rm \sim 100$ [21–24].

As we mentioned in the introduction, the Batchelor analogy of magnetic field lines and vorticity lines would hold for $\nu = \eta$ if the initial magnetic field were proportional to the vorticity field. In this special case, magnetic energy would not be amplified. In the model we investigate there are infinitely many such special initial conditions; any magnetic correlator $H_L(r)$ lacking the growing eigenfunctions in its expansion will not be amplified. This, of course, does not mean that dynamo does not exist.

We also note that the Schrödinger-type equation of the form (7) was considered for the magnetic field correlator in the case of a general velocity field in [10]. In this sense the applicability of the equation may not be restricted to the Gaussian, short-time correlated velocity field only. For example, we also investigated the modified Kazantsev model, with a more realistic, finite-time correlated velocity field. The velocity correlation time at a given wave number, k , was chosen to be of the order of the eddy turnover time at the corresponding scale. This model could not be solved analytically, however, its numerical integration gave qualitatively the same results as (1,2). We will report these results elsewhere.

We would like to thank S. C. Cowley, R. Rosner, A.

Schekochihin, and S. I. Vainshtein for many useful discussions. This work was supported by the NSF Center for magnetic self-organization in astrophysical and laboratory plasmas at the University of Chicago. S.B. would like to thank the Aspen Center for Physics, where this work was initiated.

-
- [1] L. D. Landau, E. M. Lifshitz, & L. P. Pitaevskii, *Electrodynamics of Continuous media* (Elsevier Science, Ltd. 1985).
 - [2] R. M. Kulsrud & S. W. Anderson, *ApJ* **396** (1992) 606.
 - [3] G. K. Batchelor, *Proc. Roy. Soc. London, A* **201** (1950) 405.
 - [4] R. H. Kraichnan & S. Nagarajan, *Phys. Fluids* **10** (1967) 859.
 - [5] A. P. Kazantsev, *Soviet Phys. – JETP* **26** (1968) 1031.
 - [6] A. Schekochihin, S. Boldyrev, & R. M. Kulsrud, *ApJ* **567** (2002) 828; S. Boldyrev, *ApJ* **562** (2001) 1081.
 - [7] A. Gruzinov, S. Cowley, & R. Sudan, *Phys. Rev. Lett.* **77** (1996) 4342.
 - [8] M. Chertkov, G. Falkovich, I. Kolokolov, & M. Vergasola, *Phys. Rev. Lett.*, **83** (1999) 4065.
 - [9] J. Cho, A. Lazarian, & E. T. Vishniac, *ApJ* **595** (2003) 812.
 - [10] S. I. Vainshtein & L. L. Kichatinov, *J. Fluid Mech.* **168** (1986) 73.
 - [11] A. Tsinober & B. Galanti, *Physics of Fluids* **15** (2003) 3514.
 - [12] K. Ohkitani, *Phys. Rev. E* **65** (2002) 046304.
 - [13] F. Cattaneo, in *Modelling of Stellar Atmospheres*, ed. by N. E. Peskunov, W. W. Wies, & D. F. Gray (Astron. Soc. Pac. Conf. Ser.), v. S-210 (2002).
 - [14] U. Christensen, P. Olson, & G. A. Glatzmaier, *Geophys. J. Int.* **138** (1999) 393.
 - [15] A. Schekochihin, S. C. Cowley, J. L. Maron, & J. C. McWilliams, *astro-ph/0308336*.
 - [16] T. A. Yousef, A. Brandenburg, & G. Rüdiger, *astro-ph/0302425*.
 - [17] R. H. Kraichnan, *Phys. Fluids*, **11** (1968) 945.
 - [18] L. D. Landau & E. M. Lifshitz, *Quantum mechanics* (Elsevier Science, Ltd. 1985).
 - [19] G. S. Golitsyn, *Sov. Phys. Doklady*, **5** (1960) 536.
 - [20] K. Moffatt, *J. of Fluid Mechanics*, **11**, 2 (1961) 625.
 - [21] A. Gailitis et al, *Phys. Rev. Lett.* **86** (2001) 3024.
 - [22] A. Gailitis, O. Lielausis, E. Platācis, G. Gerbeth, & F. Stefani, *Rev. Mod. Phys.* **74** (2002) 973.
 - [23] M. Bourgoin, et al, *Phys. Fluids*, **14** (2002) 3046.
 - [24] C. B. Forest, et al, *Magnetohydrodynamics* **38** (2002) 107.